


SMM C/P 1988 DOY156 75 +/- 0 degrees



tstart=DOY156 1: 0 +/- 0.0 [min.]
radial velocity= 421. +/- 0. [km/sec]

This is a 1 order run

INPUT:

| year | day | time | radii | error in radius | angle | error in angle | datatype |
|------|-----|---------|-------|-----------------|-------|----------------|----------|
| 1988 | 156 | 1:30: 6 | 2.10 | +/- 0.15 | 75.0 | +/- 0.5 | SMM |
| 1988 | 156 | 2: 3: 9 | 3.30 | +/- 0.15 | 75.0 | +/- 1.0 | SMM |

Structure being measured:
Top of Mound

OUTPUT:

START TIME:

Event began at $t = 0 + -0.302E+02 +/- 0.000E+00$ [minutes]

In more convenient units:

Event began at DOY 156 1:00 +/- 0.000E+00 [minutes]

Event began at DOY 156 1: 0 +/- 0.000E+00 [minutes]

Origin defined at DOY156 1:30

RADIAL INFORMATION:

RADIAL VELOCITY = 421. +/- 0. km/sec

The computed RADIAL coefficients and errors FROM NLLST2 are:

$c(1) = 0.210E+01 +/- 0.239E-06$ [solar radii]
 $c(2) = 0.363E-01 +/- 0.102E-07$ [radii/min]
 $c(2) = 0.421E+03 +/- 0.118E-03$ [km/sec]
 $c(3) = 0.000E+00 +/- 0.000E+00$ [radii/min**2]
 $c(3) = 0.000E+00 +/- 0.000E+00$ [km/sec**2]

ANGULAR INFORMATION:

ANGULAR VELOCITY = 0.0000E+00 +/- 0.0000E+00[radians/sec]

The computed ANGULAR coefficients and errors FROM NLLST2 are:

$a(1) = 75. +/- 0.$ [degrees]
 $a(2) = 0.000E+00 +/- 0.000E+00$ [degrees/min]
 $a(2) = 0.000E+00 +/- 0.000E+00$ [rad/sec]

a(3) = 0.000E+00 +/- 0.000E+00 [degrees/min**2]
a(3) = 0.000E+00 +/- 0.000E+00 [rad/sec**2]

THE TANGENTIAL AND TOTAL VELOCITIES AT EACH DATA POINT ARE:

The TANGENTIAL velocity at 1:30 = 0. km/sec

The TOTAL velocity at 1:30 = 421. km/sec

The TANGENTIAL velocity at 2: 3 = 0. km/sec

The TOTAL velocity at 2: 3 = 421. km/sec

The following info is for estimating goodness of the fit

Number data points = number of coefficients. No error analysis is possible.

Correlation coefficient is not computable!