

## Empirical Geomagnetic Inputs to the TIE-GCM

Stan Solomon, 3/18/16

The hierarchy of auroral/geomagnetic control parameters in the TIE-GCM is somewhat confusing, because there are so many options.

First of all, there is a switch called `POTENTIAL_MODEL` which can be set to either "HEELIS" (the default) or "WEIMER".

In either case, the model may then be run with namelist input parameters specifying the inputs to the potential model, or with time-dependent parameters read from a file.

For Heelis runs: the model will use the cross-tail potential `CTPOTEN` and hemispheric power `POWER` specified in the namelist inputs, if present. Alternatively, it can use `KP` specified in the namelist inputs, from which `CTPOTEN` and `POWER` are calculated using correlation relationships. These can also be time-dependent arrays in the namelist inputs. If `CTPOTEN`, `POWER`, and `KP` are **\*not\*** present, the model reads a geophysical index file, specified by `GPI_NCFILE`, from which it gets time-dependent 3-hour `KP`. A GPI file covering 1960-2015 is provided with the model download.

For Weimer runs: the model will use interplanetary magnetic field and solar wind parameters `BYIMF`, `BZIMF`, `SWDEN`, and `SWVEL` specified by the namelist inputs. If these are not present, it will read them from a file specified by `IMF_FILE`. A GPI file can also be specified for Weimer runs, but only for solar irradiance inputs. Selected 1-year IMF files are provided with the model download, and others are available at <http://download.hao.ucar.edu/pub/tgcm/>.

Solar irradiance input control is similar. The model uses `F107` and `F107A` if specified in the namelist inputs, and otherwise gets them from the GPI file. It is possible to mix solar and auroral inputs between namelist and file-based, for instance, you could run with time-dependent `KP` from the GPI file but constant `F10.7` and `F10.7` from namelist inputs.

So, if you're trying to do a "Heelis/GPI" run, or a "Weimer/IMF" run (in our vernacular), don't set namelist inputs because they will override the file-based inputs. For more details, see chapter 5 of the user guide, and the extensive comment in subroutine `inp_solar` in file `input.F`.

This note does not address more sophisticated auroral/geomagnetic specifications such as AMIE and CMIT.